

**ASUHAN KEPERAWATAN MEDIKAL BEDAH DENGAN GANGGUAN  
SISTEM SARAF STROKE NON HEMORAGIC DENGAN MASALAH  
KEPERAWATAN PERFUSI SELEBRAL TIDAK EFEKTIF PADA  
TN YA DI BANGSAL ANNA DI RUMAH SAKIT  
SUAKA INSAN**

*Marianus Aloysio<sup>1</sup> , Lucia Andi Chrimilasari<sup>2</sup>*

**INTISARI**

**Latar belakang:** Stroke merupakan penyebab kematian ketiga di negara industri setelah penyakit arteri koroner (13%) dan kanker (12%). Prevalensi stroke bervariasi di berbagai belahan dunia

**Tujuan:** menggambarkan proses dan hasil asuhan keperawatan medikal bedah dalam mengatasi masalah risiko perfusi jaringan selebral tidak efektif dengan mobilisasi progresif.

**Hasil:** tindakan mobilisasi progresif

**Kesimpulan:** setelah dilakukan implementasi pemberian mobilisasi progresif pada pasien teratasi ditandai dengan tercapainya kriteria hasil ekstermitas pasien membaik, meringis berkurang, mampu melakukan penanganan mobilisasi secara mandiri.

**Kata kunci:** risiko perfusi jaringan selebral tidak efektif, mobilisasi progresif, stroke non hemoragic

---

---

*STIKES Suaka Insan Banjarmasin*

---

<sup>1</sup> Mahasiswa Sekolah Tinggi Ilmu Kesehatan Suaka Insan Banjarmasin

<sup>2</sup> Dosen Sekolah Tinggi Ilmu Kesehatan Suaka Insan Banjarmasin

**MEDICAL SURGICAL NURSING CARE FOR NERVOUS SYSTEM  
DISORDERS NON-HEMORRHAGIC STROKE WITH INEFFECTIVE  
CELEBRAL PERFUSION TREATMENT PROBLEMS IN TN YA IN  
ANNA ward AT INSAN SUAKA HOSPITAL**

*Marianus Aloysio<sup>3</sup>, Lucia Andi Chrimilasari<sup>4</sup>*

**ABSTRACT**

**Introduction:** Stroke is the third cause of death in industrialized countries after coronary artery disease (13%) and cancer (12%). The prevalence of stroke varies in different parts of the world

**Purpose:** describes the process and results of medical surgical nursing care in overcoming the risk of ineffective cerebral tissue perfusion with progressive mobilization..

**Results:** progressive mobilization actions

**Conclusion:** After the implementation of progressive mobilization, the patient resolved, as indicated by the achievement of the patient's extremity outcome criteria, improved, grimacing reduced, and was able to handle mobilization independently.

**Keywords:** risk of ineffective cerebral tissue perfusion, progressive mobilization, non-hemorrhagic stroke

---

---

*STIKES Suaka Insan Banjarmasin*

---

<sup>3</sup> The student of program Suaka Insan School of health sciences higher education

<sup>4</sup> The lecture of program Suaka Insan School of health sciences higher education